



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Journal of Mycology

VOLUME 10—MARCH 1904

MORGAN—A New Melogramma	49
COCKERELL—Some Fungi Collected in New Mexico	49
DUDLEY AND THOMPSON—Notes on Californian Uredineæ and Descriptions of New Species.....	52
KELLERMAN—Ohio Fungi, Fascicle IX	55
KELLERMAN—Minor Mycological Notes. III.....	62
KELLERMAN—Index to Uredineous Culture Experiments, Concluded	64
KELLERMAN—Notes from Mycological Literature. IX.....	81
KELLERMAN—Elementary Mycology.....	90
EDITOR'S NOTES.....	96

A NEW MELOGRAMMA.

A. P. MORGAN.

MELOGRAMMA PATENS Morgan n. sp.—Stroma depressed-pulvinate, cinereous then black, the surface mammillate by the numerous subjacent perithecia. Perithecia, small, ovoid, papillate, lying in a single layer. Asci cylindric, with a short stalk, octosporous, paraphysate, 120-130 x 7-9 mic.; spores fusiform, slightly curved, 3-septate, yellow-brown, 18-22 x 6-7 mic.

Growing on dead branches of *Carpinus americana*, Preston, Ohio, 1896. Stromata gregarious, seated on the inner bark and erumpent through the periderm, 2-3 mm. in diameter, containing from 50 to more than 100 perithecia.

The nearest relative to this species appears to be *Melogramma caucasicus* Jaczewski. The common *Melogramma vagans* DeNot. grew along with it on the same branches, sometimes mingled together, but the difference between them is perfectly obvious to the naked eye.

SOME FUNGI COLLECTED IN NEW MEXICO.

T. D. A. COCKERELL.

For several years I have been preparing a catalogue of the fungi of New Mexico; but it is still hardly more than a fragment, and I do not suppose I shall ever be able to make it anything better, now that I have moved to Colorado. It contains, however, some records which seem worth publishing, as additions